

Literaturverzeichnis zur Veröffentlichung:

SOMMER, R., ZIARNETZKY, V., MESSLINGER, U., ZAHNER, V. (2019): Der Einfluss des Bibers auf die Artenvielfalt semiaquatischer Lebensräume – Sachstand und Metaanalyse für Europa und Nordamerika. Naturschutz und Landschaftsplanung 51 (3), 108-115.

- ANDERSON, C. B., ROSEMOND, A. D. (2007): Ecosystem engineering by invasive exotic beavers reduces in-stream diversity and enhances ecosystem function in Cape Horn, Chile. *Oecologia*, 154, 141–153.
- ANDERSON, N. L., PASZKOWSKI, C. A., HOOD, G. A. (2014): Linking aquatic and terrestrial environments: can beaver canals serve as movement corridors for pond-breeding amphibians? *Animal Conservation* 18, 287-294.
- ANDRZEJEWSKA-WIERZBICKA, M., BERESZYŃSKI, M. (2000): Ekologia oraz środowiskotwórcza rola bobra europejskiego (*Castor fiber* Linnaeus, 1758) w dolinie strumienia Kończak. (Ecology and environmental role of the European beaver in the Kończak stream valley). *Roczniki AR w Poznaniu, Zootechnika* 52, 3-24.
- ARNDT, E., DOMDEI, J. (2011): Influence of beaver ponds on the macroinvertebrate benthic community in lowland brooks. *Polish Journal of Ecology* 59, 799-811.
- AZNAR, J. C., DESROCHERS, A. (2008): Building for the future: abandoned beaver ponds promote bird diversity. *Ecoscience* 15, 250-257.
- BAKER, B. W., DUCHARME, H. C., MITCHELL, D. C. S., STANLEY, T. R., PEINETTI, H. R. (2005): Interaction of beaver and elk herbivory reduces standing crop of willow. *Ecological Applications* 15, 110-118.
- BEARD, E. B. (1953): The Importance of Beaver in Waterfowl Management at the Seney National Wildlife Refuge. *Journal of Wildlife Management* 17, 398-436.
- BALAČIAUSKAS, L., BALAČIAUSKIENĖ, L., TRAKIMAS, G. (2001): Beaver influence on amphibian breeding in the agrolandscape. In: CZECH, A., SCHWAB, G. (Hrsg.), *The European Beaver in a new millennium. Proceedings of 2nd European Beaver Symposium, 2000 Białowieża, Poland*. Carpathian Heritage Society, Kraków, 105-112.
- BARTEL, R. A., HADDAD, N. M., WRIGHT, J. P. (2010): Ecosystem engineers maintain a rare species of butterfly and increase plant diversity. *Oikos* 119, 883-890.
- CIECHANOWSKI, M., KUBIC, W., RYMKIEWICZ, A., ZWOLICKI, A. (2011): Reintroduction of beavers *Castor fiber* may improve habitat quality for vespertilionid bats foraging in small river valleys. *European Journal of Wildlife Research* 57, 737-747.
- COLLEN, P., GIBSON R. J. (2001): The general ecology of beavers (*Castor* spp.), as related to their influence on stream ecosystems and riparian habitats, and the subsequent effects on Fish – a review. *Reviews in Fish Biology and Fisheries* 10, 439–461.
- CUNNINGHAM, J. M., CALHOUN, A. J. K., GLANZ, W. E. (2007): Pond-breeding amphibian species richness and habitat selection in a beaver-modified landscape. *Journal of Wildlife Management* 71, 2517–2526.
- CZERNIAWSKI, R., SŁUGOCKI, Ł, KOWALSKA-GÓRALSKA, M. (2017): Effects of beaver dams on the zooplankton assemblages in four temperate lowland streams (NW Poland). *Biologia* 72, 417-430.
- DALBECK, L. (2011): Biberlichtungen als Lebensraum für Heuschrecken in Wäldern der Eifel. *Articulata* 26, 97-108.
- , JANSSEN, J., VÖLSGEN, S. L. (2014): Beavers (*Castor fiber*) increase habitat availability, heterogeneity and connectivity for common frogs (*Rana temporaria*). *Amphibia- Reptilia* 35, 321-329.
- , LÜSCHER, B., OHLHOFF, D. (2007): Beaver ponds as habitat of amphibian communities in a central European highland. *Amphibia-Reptilia* 28, 493-501.
- DERWICH, A., BRZUSKI, P., HĘDRZAK, M. (2007): *Bóbr w biotopach Bieszczadów Wysokich* (Beavers in biotopes of the western Bieszczady). Wydawnictwo Drukpol, Kraków, 112 pp.
- DEVAUX, B. (2000): Der Schutz der Europäischen Sumpfschildkröte *Emys orbicularis* (L.) in der Provence (Frankreich). *Stapfia* 69, 195-204.

- ECKE, F., LEVANONI, O., AUDET, J., CARLSON, P., EKLÖF, K., HARTMAN, G., MCKIE, B., LEDESMA, J., SEGERSTEN, J., TRUCHY, A., FUTTER, M. (2017): Meta-Analysis of environmental effects of beaver in relation to artificial dams. *Environmental Research Letters* 12, 113002.
- FRANCE, R. L. (1997): The importance of beaver lodges in structuring littoral communities in boreal headwater lakes. *Canadian Journal of Zoology* 75, 1009-1013.
- FREITAG, H., STUBBE, M., HEIDECHE, D. (2001): Das Makrozoobenthos in der Zönosestruktur und die Saprobie unter Einfluss des Elbe-Bibers. *Säugetierkundliche Informationen* 5, 35-56.
- FYODOROV, F. V., YAKIMOVA, A. E. (2012): Changes in Ecosystems of the Middle Taiga due to the impact of Beaver Activities, Karelia, Russia. *Baltic Forestry* 18, 278-287.
- GROVER, A. M., BALDASSARRE, G. A. (1995): Bird species richness within beaver ponds in South-Central New York. *Wetlands* 15, 108-118.
- GURNELL, A. M. (1998): The hydromorphological effects of beaver dam-building activity. *Progress in Physical Geography* 22, 167-189.
- GURNELL, J., GURNELL, A. M., DEMERITT, D., LURZ, P. W. W., SHIRLEY, M. D. F., RUSHTON, S. P., FAULKES, C. G., NORBERT, S., HARE, E. J. (2009): The feasibility and acceptability of reintroducing the European beaver to England. Report for: Natural England and the People's Trust for Endangered Species.
- HANÖFFER, S., SCHURLI, C. (2003): Einfluss der Dammbauaktivität des Bibers auf die Fischzönosen im Mühlbach bei Niederhummel (Lkr. Freising). Diplomarbeit Hochschule Weihenstephan, Freising. VI, 123 S.
- HÄGGLUND, A., SJÖBERG, G. (1999): Effects of beaver dams on the fish fauna of forest streams. *Forest Ecology and Management*, 115, 259-266.
- HARTHUN, M. (1999): Der Einfluss des Bibers (*Castor fiber albicus*) auf die Fauna (Odonata, Mollusca, Trichoptera, Ephemeroptera, Diptera) von Mittelgebirgsbächen in Hessen (Deutschland). *Limnologica* 29, 449-464.
- HOOD, G. A., BAYLEY, S. E. (2009): A comparison of riparian plant community response to herbivory by beavers (*Castor canadensis*) and ungulates in Canada's boreal mixed-wood forest. *Forest Ecology and Management* 258, 1979-1989.
- , LARSSON, D. G. (2014): Beaver-created habitat heterogeneity influences aquatic invertebrate assemblages in Boreal Canada. *Wetlands* 34, 19-29.
- , LARSON, D. G. (2015): Ecological engineering and aquatic connectivity: a new perspective from beaver-modified wetlands. *Freshwater Biology* 60, 198-208.
- JANISZEWSKI, P., HANZAL, V., MISIUKIEWICZ, W. (2014): The eurasian beaver (*Castor fiber*) as a key stone species – a literature review. *Baltic Forestry* 20, 277-286.
- JONES, C. G., LAWTON, J. H., SHACHAK, M. (1994): Organisms as ecosystem engineers. *Oikos* 69, 373-386.
- KARRAKER, N. E., GIBBS, J. P. (2009): Amphibian production in forested landscapes in relation to wetland hydroperiod: a case study of vernal pools and beaver ponds. *Biological Conservation* 142, 2293–2302.
- KEMP, P. S., WORTHINGTON, T. A., LANGFORD, T. E. L., TREE, A. R. J., GAYWOOD, M. J. (2012): Qualitative and quantitative effects of reintroduced beavers on stream fish. *Fish and Fisheries* 13, 158-181.
- KESMINAS, V., STEPONĖNAS, A., PLIŪRAITĖ, V., VIRBICKAS, T. (2013): Ecological Impact of Eurasian Beaver (*Castor fiber*) activity on fish communities in Lithuanian trout streams. *Annual Set The Environment Protection Rocznik Ochrona Środowiska* 15, 59-80.
- KUKUŁA, K., BYLAK, A. (2010): Ichthyofauna of a mountain stream dammed by beaver. *Archives of Polish Fisheries* 18, 33-43.
- LAW, A., MCLEAN, F., WILLBY, N. J. (2016): Habitat engineering by beaver benefits aquatic biodiversity and ecosystem processes in agricultural streams. *Freshwater Biology* 61, 486-499.
- , GAYWOOD, M. J., JONES, K. C., RAMSAY, P., WILLBY, N. J. (2017): Using ecosystem engineers as tool in habitat restoration and rewilding: beaver and wetlands. *Science of the Total Environment* 605-606, 1021-1030.
- METTS, B. S., LANHAM, J. D., RUSSELL, K. R. (2001): Evaluation of Herpetofaunal Communities on Upland Streams and Beaver-Impounded Streams in the Upper Piedmont of South Carolina. *The American Midland Naturalist* 145, 54-65.

- NAIMAN, R. J., MELILLO, J. M., HOBBIE, J. E. (1986): Ecosystem alteration of boreal forest streams by beaver (*Castor canadensis*). *Ecology* 67, 1254–1269.
- NUMMI, P. (1989): Simulated effects of the beaver on vegetation, invertebrates and ducks. *Annales Zoologici Fennici* 26, 43-52.
- (1992): The importance of beaver ponds to waterfowl broods: an experiment and natural tests. *Annales Zoologici Fennici* 29, 47-55.
- , HAHTOLA, A. (2008): The beaver as an ecosystem engineer facilitates teal breeding. *Ecography* 31, 519-524.
- , HOLOPAINEN, S. (2014): Whole-community facilitation by beaver: ecosystem engineer increases waterbird diversity. *Aquatic Conservation: Marine and freshwater Ecosystems* 24, 623-633.
- , PÖYSÄ, H. (1997): Population and community level responses in *Anas*-species to patch disturbance caused by an ecosystem engineer, the beaver. *Ecography* 20, 580-584.
- , KATTAINEN, S., ULANDER, P., HAHTOLA, A. (2011): Bats benefit from beavers: a facilitative link between aquatic and terrestrial food webs. *Biodiversity and Conservation* 20, 851–859.
- PARKER, H., RØNNING, O. C. (2007): Low potential for restraint of anadromous salmonid reproduction by beaver *Castor fiber* in the Numedalslgen river catchment, Norway. *River Research and Applications* 23, 752-762.
- PLIŪRAITĖ, V., KESMINAS, V. (2012): Ecological impact of Eurasian beaver (*Castor fiber*) activity on macroinvertebrate communities in Lithuanian trout streams. *Central European Journal of Biology* 7, 101-114.
- POPESCU, V. D., GIBBS, J. P. (2009): Interactions between climate, beaver activity, and pond occupancy by the cold-adapted mink frog in New York State, USA. *Biological Conservation* 142, 2059-2068.
- ROLLAUFS, P., HERING, D., LOHSE, S. (2001): Composition, invertebrate community and productivity of a beaver dam in comparison to other stream habitat types. *Hydrobiologia* 459, 201-212.
- ROSELL, F., BOSZER, O., COLLEN, P., PARKER, H. (2005): Ecological impact of beavers *Castor fiber* and *Castor canadensis* and their ability to modify ecosystems. *Mammal Review* 35, 248-276.
- RUNYON, M. J., TYERS, D. B., SOWELL, B. F., GOWER, C. N. (2014): Aspen restoration using beaver on the Northern Yellowstone winter range under reduced ungulate herbivory. *Restoration Ecology* 22, 555-561.
- RUSSELL, K. R., MOORMAN, C. E., EDWARDS, J. K., METTS, B. S., GUYNN, D. C. JR. (1999): Amphibian and reptile communities associated with beaver (*Castor canadensis*) ponds and unimpounded streams in the Piedmont of South Carolina. *Journal of Freshwater Ecology* 14, 149–158.
- SAMAS, A. (2016): Impact of the keystone species, the eurasian beaver (*Castor fiber*), on habitat structures and its significance to mammals. PhD thesis, Vilnius University, 133 pp.
- SCHLOEMER, S., DALBECK, L. (2014): Der Einfluss des Europäischen Bibers (*Castor fiber*) auf Mittelgebirgsbäche der Nordeifel (NRW) am Beispiel der Libellenfauna (Odonata). In: NITSCHKE, K. A. (Hrsg.), Tagungsband Ergebnisse der Nationalen Bibertagung in Dessau-Roßlau, Sachsen-Anhalt, pp. 25-29.
- SIDOROVICH, V. (2011): Analysis of vertebrate predator-prey community. Tesey Publisher, Minsk, 736 pp.
- SIKORA, A., RYS, A. (2004): Distribution, abundance and habitat preferences of the white-backed woodpecker *Dendrocopos leucotos* in the regions of Warmia and Masuria. *Notatki Ornitologiczne* 45, 253–262.
- SMITH, J. M., MATHER, M. E. (2013): Beaver dams maintain fish biodiversity by increasing habitat heterogeneity throughout a low-gradient stream network. *Freshwater Biology* 58, 1523-1538.
- SNODGRASS, J. W., MEFFE, G. K. (1998): Influence of beavers on stream fish assemblages: effects of pond age and watershed position. *Ecology* 79, 928-942.
- STEVENS, C. E., PASZKOWSKI, C. A., FOOTE, A. L. (2007): Beaver (*Castor canadensis*) as a surrogate species for conserving anuran amphibians on boreal streams in Alberta, Canada. *Biological Conservation* 134, 1–13.
- STRINGER, A. P., GAYWOOD, M. J. (2016): The impact of beavers *Castor* ssp. on biodiversity and the ecological basis for their reintroduction to Scotland, UK. *Mammal Review* 46, 270-283.

- TÖRNBLOM, J., ANGELSTAM, P., HARTMAN, G., HENRIKSON, L., SJÖBERG, G. (2011): Towards a research agenda for water policy implementation: knowledge about beaver (*Castor fiber*) as a tool for water management with a catchment perspective. *Baltic Forestry* 17, 154-161.
- TUMIEL, T. (2008): Abundance and distribution of the three-toed woodpecker in the Puszcza Knyszynska forest in 2005-2007. *Notatki Ornitologiczne* 49, 74-80.
- WRIGHT, J. P. (2009): Linking populations to landscapes: richness scenarios resulting from changes in the dynamics of an ecosystem engineer. *Ecology* 90, 3418-3429.
- , JONES, C. G., FLECKER, A. S. (2002): An ecosystem engineer, the beaver, increases species richness at the landscape scale. *Oecologia* 132, 96-101.
- ZAHNER, V., HANÖFFER, S., SCHURLI, C., MÜLLER, S. (2006): Beaver induced structure change along a stream in Bavaria and its influence on fish fauna and indicator beetles. In: ZAHNER, V., SCHWAB, G., SCHMIDBAUER, M., NITSCHKE, K.-A., BUSER, P. E. (Hrsg.): Abstracts from the 4th European Beaver Symposium. Fachhochschule Weihenstephan, Freising, Germany, pp. 68.

Literaturquellen, die ausschließlich für die Meta-Analyse genutzt wurden

- ANDERSON, C. B., GRIFFITH, C. R., ROSEMOND, A. D., ROZZI, R., DOLLENZ, O. (2006): The effects of invasive North American beavers on riparian plant communities in Cape Horn, Chile. Do exotic beavers engineer differently in sub-Antarctic ecosystems? *Biological Conservation* 128, 467-474.
- BROMLEY, C. K., HOOD, G. A. (2013): Beavers (*Castor canadensis*) facilitate early access by Canada geese (*Branta canadensis*) to nesting habitat and areas of open water in Canada's boreal wetlands. *Mammalian Biology* 78, 73-77.
- BRZYSKI, J. R., SCHULTE, B. A. (2009): Beaver (*Castor canadensis*) Impacts on Herbaceous and Woody Vegetation in Southeastern Georgia. *The American Midland Naturalist* 162, 74-86.
- CLIFFORD, H. F., WILEY, G. M., CASEY, R. J. (1993): Macroinvertebrates of a beaver-altered boreal stream of Alberta, Canada, with special reference to the fauna on the dams. *Canadian Journal of Zoology* 71, 1439-1447.
- CRISLER, J. D., RUSSELL, F. L. (2010): Patterns in Beaver Herbivory in South-Central Kansas Riparian Woodlands. *Transactions of the Kansas Academy of Science*, 113, 161-176.
- DONKOR, N. T., FRYXELL, J. M. (1999): Impact of beaver foraging on structure of lowland boreal forests of Algonquin Provincial Park, Ontario. *Forest Ecology and Management* 118, 83-92.
- KRYLOV, A. V. (2008): Impact of the Activities of Beaver on the Zooplankton of a Piedmont River (Mongolia). *Inland Water Biology* 1, 73-75.
- LAW, A., JONES, K. C., WILLBY N. J. (2014): Medium vs. Short-term effects of herbivory by Eurasian beaver on aquatic vegetation. *Aquatic Botany* 116, 27-34.
- LESICA, P., MILES, S. (2004): Beavers indirectly enhance the growth of russian olive and tamarisk along eastern montana rivers. *Western North American Naturalist* 64, 93-100.
- MARGOLIS, B. E., RAESLY, R. L., SHUMWAY, D. L. (2001): The effect of beaver created wetlands on the benthic macroinvertebrate assemblages of two Appalachian streams. *Wetlands* 21, 554-563.
- MARTINSEN, G. D., DRIEBE, E. M., WHITHAM, T. G. (1998): Indirect interactions mediated by changing plant chemistry: Beaver browsing benefits beetles. *Ecology* 79, 192-200.
- MCDOWEL, D. M., NAIMAN, R. J. (1986): Structure and function of a benthic invertebrate stream community as influenced by beaver (*Castor canadensis*). *Oecologia* 68, 481-489.
- MITCHELL, C. C., NIERING, W. A. (1993): Vegetation change in a topogenic bog following beaver flooding. *Bulletin of the Torrey Botanical Club* 120, 136-147.
- NAIMAN, R. J., MCDOWELL, D. M., FARR, B. S. (1984): The influence of beaver (*Castor canadensis*) on the production dynamics of aquatic insects. *Verhandlungen des Internationalen Verein Limnologie* 22, 1801-1810.
- NOLET, B., HOEKSTRA, A., OTTENHEIM, M. M. (1994): Selective foraging on woody species by the beaver *Castor fiber*, and its impact on a riparian willow forest. *Biological Conservation* 70, 117-128.

- OBIDZIŃSKI, A., ORCZEWSKA, A., CIELOSZCZYK, P. (2011): The impact of beavers' (*Castor fiber* L.) lodges on vascular plant species diversity in forest landscape. *Polish Journal of Ecology* 59, 63-73.
- SCHLOSSER, I. (1995): Dispersal, boundary processes, and trophic-level interactions in streams adjacent to beaver ponds. *Ecology* 76, 908-925.
- SUZUKI, N., MCCOMB, B. (2004): Associations of small mammals and amphibians with beaver-occupied streams in the Oregon Coast Range. *Northwest Science* 78, 286-292.
- WESTBROOK, C. J., COOPER, D. J., BAKER, B. W. (2011): Beaver assisted river valley formation. *River Research and Applications* 27, 247-256.
- WRIGHT, J. P., FLECKER, A. S., JONES, C. G. (2003): Local vs. Landscape controls on plant species richness in beaver meadows. *Ecology* 84, 3162-3173.